



INVESTOR IN PEOPLE

The Patent Office  
Concept House  
Cardiff Road  
Newport  
South Wales  
NP10 8QQ

## PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

REC'D 24 AUG 2004

WIPO

PCT

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.


In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.

Signed

Dated

16 August 2004




The  
Patent  
Office

25 JUL 2003 E825263-3 002944  
P0147700 0-00-0317393.7

# Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)



The Patent Office

Cardiff Road  
Newport  
Gwent NP9 1RH

1. Your reference SMC 60613/GB/P1

2. Patent application number  
(The Patent Office will fill in this part) 0317393.7 25 JUL 2003

3. Full name, address and postcode of the or of each applicant (underline all surnames) Avecia Limited  
Hexagon House  
Blackley  
Manchester, M9 8ZS

Patents ADP number (if you know it)

If the applicant is a corporate body, give the country/state of its incorporation

07764137001

United Kingdom

4. Title of the invention PROCESS AND COMPOUNDS

5. Name of your agent (if you have one) REVELL, Christopher

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

Avecia Limited  
Hexagon House  
PO Box 42  
Blackley  
Manchester M9 8ZS

Patents ADP number (if you know it)

6969885001

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number	Country	Priority application number (if you know it)	Date of filing (day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application	Number of earlier application	Date of filing (day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:

- a) any applicant named in part 3 is not an inventor, or
- b) there is an inventor who is not named as an applicant, or
- c) any named applicant is a corporate body.

See note (d))

APPLICANTS

AVECIA LIMITED

TITLE


PROCESS AND COMPOUNDS

---

PROCESS AND COMPOUNDS

The present invention concerns a process and intermediate compounds useful in the preparation of statins, particularly atorvastatin.

According to the present invention, there is provided a process for the preparation of a compound of formula (7):



wherein

R<sup>1</sup> represents an alkyl group, such as a C<sub>1-6</sub> alkyl group, and preferably an isopropyl group;


R<sup>2</sup> represents an aryl group, preferably a phenyl group

R<sup>3</sup> represents an aryl group, preferably a 4-fluorophenyl group


X represents a group of formula -COZ, wherein Z represents -OR<sup>4</sup>, in which R<sup>4</sup> represents an alkyl, preferably a methyl or ethyl, group, or -NR<sup>5</sup>R<sup>6</sup>, wherein R<sup>5</sup> and R<sup>6</sup> each independently represent H, alkyl, or aryl, and preferably R<sup>5</sup> is H and R<sup>6</sup> is phenyl

which comprises


a) cyanating a compound of formula (1):




wherein Y represents a halo group, preferably Cl or Br; P<sup>1</sup> represents a protecting group, and W represents =O or -OP<sup>2</sup>, in which P<sup>2</sup> represents a protecting group, to give a compound of formula (2):




b) reducing the compound of formula (2) to give a compound of formula (3):




c) coupling the compound of formula (3) with a compound of formula (4):



to give a compound of formula (5):




d) when W represents  $-OP^2$ , deprotecting and then oxidising the compound of formula (5) to give a compound of formula (6):



and

e) subjecting the compound of formula (5) when W represents  $=O$ , or compound of formula (6) to ring-opening, and removal of any remaining protecting groups, to give a compound of formula (7):



Protecting groups which may be represented by  $P^1$  and  $P^2$  include alcohol protecting groups, examples of which are well known in the art. Particular examples include tetrahydropyranyl and benzyl groups. Preferred protecting groups are silyl groups, for example triaryl- and especially trialkylsilyl groups. Especially preferred examples are trimethylsilyl and t-butyl dimethylsilyl groups.

Cyanation of compounds of formula (1) can be achieved by methods known in the art for displacing a halo group with a cyanide. Preferably, the process comprises contacting the compound of formula (1) with a source of cyanide. Preferred sources of cyanide include cyanide salts, especially ammonium or alkali metal cyanides, particularly sodium or potassium cyanide. A particularly preferred process comprises contacting the compound of formula (1) with 5 molar equivalents of KCN in the presence of dimethylsulfoxide solvent at a temperature of, for example, about  $50^\circ\text{C}$ .

Reduction of compounds of formula (2) can be achieved using reduction systems known in the art for the reduction of nitrile groups. Preferred examples include reduction

with Raney nickel or with hydrogen in the presence of a catalyst, such as palladium on carbon. When palladium on carbon catalysed hydrogenation is employed, preferred conditions comprise the use of methanol solvent at elevated temperature, such as about 40°C, in the presence of from about 0.01 to 100 molar equivalents of ammonia.

The coupling of the compound of formula (3) with the compound of formula (4) may employ conditions analogous to those given in WO89/07598 for the corresponding coupling. The conditions preferably comprise refluxing the compounds of formula (3) and (4) in a hydrocarbon solvent, such as toluene or cyclohexane, or mixtures thereof, followed by contact with aqueous acid, such as aqueous HCl.

When W represents  $OP^2$ , the protecting group may be removed to form a hydroxy group by methods known in the art for the removal of the given protecting group. For example, silyl protecting groups may be removed by contact with a source of fluoride ion, such as tetrabutylammonium fluoride.

Oxidation of compounds formed by deprotection of compounds wherein W represents  $-OP^2$  may employ conditions known in the art for the oxidation of pyranols to pyranones, and include those given in "Comprehensive Organic Transformations", R.C. Larock, 2<sup>nd</sup> Ed (1999) p 1670, published by Wiley VCH, incorporated herein by reference. Preferred oxidation systems include  $Ag_2CO_3$ /Celite, especially Celite J2, or bromine.


Ring opening of the compounds of formula (5), when W represent =O or formula (6) may employ conditions known in the art for ring opening of a pyranone. Preferably, the ring is opened by contact with a base, such as sodium hydroxide. Methanol is conveniently employed as solvent.

Remaining protecting groups may be removed by methods known in the art for the removal of the given protecting group. For example, silyl protecting groups may be removed by contact with a source of fluoride ion, such as tetrabutylammonium fluoride.

It will be recognised that when X represents a group of formula  $-COOR^4$ , this may be converted to a group wherein X represents  $-CONR^5R^6$  at any stage during the process, for example by reaction of the corresponding compounds of formulae (2), (3), (4), (5), (6) or (7) with a compound of formula  $HNR^5R^6$ .


It will also be recognised that compounds of formulae (2), (3) and (4) may also be subjected to oxidation (when W represents  $-OH$ ) or deprotection and oxidation (when W represents  $-O$ -protecting group) to form the corresponding compound wherein W represents =O.

Preferred compounds of formula (1) are compounds of formula:




wherein W,  $P^1$  and Y are as previously described.

Preferred compounds of formula (2) are compounds of formula:




wherein W and P¹ are as previously described.

Preferred compounds of formula (3) are compounds of formula:




wherein W and P¹ are as previously described.

Preferred compounds of formula (5) are of formula:




wherein R¹, R², R³, W, X and P¹ are as previously described.

Preferred compounds of formula (6) are of formula:



wherein R¹, R², R³, and X are as previously described.


Preferred compounds of formula (7) are of formula:



wherein R¹, R², R³, and X are as previously described.

Compounds of formula (7) are advantageously converted to pharmaceutically acceptable salts, especially their calcium salts.

Compounds of formula (4) are advantageously prepared by the methods given in J. Med. Chem., 1991, **34**, pp357-366. Particularly preferred compounds of formula (4) are compounds of formula:




Compounds of formula (1) are advantageously prepared by enzyme catalysed condensation of acetaldehyde and 2-haloacetaldehyde, for example using the method given in US patent 5,795,749.

5 Compounds of formulae (2) and (3) and, when W is  $OP^2$ , formula (5) form further aspects of the present invention.



CLAIMS

1. A process for the preparation of a compound of formula (7):



5 wherein

R<sup>1</sup> represents an alkyl group, such as a C<sub>1-6</sub> alkyl group, and preferably an isopropyl group;


R<sup>2</sup> represents an aryl group, preferably a phenyl group

R<sup>3</sup> represents an aryl group, preferably a 4-fluorophenyl group


- 10 X a group of formula -COZ, wherein Z represents -OR<sup>4</sup>, in which R<sup>4</sup> represents an alkyl, preferably a methyl or ethyl, group, or -NR<sup>5</sup>R<sup>6</sup>, wherein R<sup>5</sup> and R<sup>6</sup> each independently represent H, alkyl, or aryl, and preferably R<sup>5</sup> is H and R<sup>6</sup> is phenyl

which comprises

- 15 a) cyanating a compound of formula (1):




wherein Y represents a halo group, preferably Cl or Br; P<sup>1</sup> represents a protecting group, and W represents =O or -OP<sup>2</sup>, in which P<sup>2</sup> represents a protecting group, to give a compound of formula (2):




20


- b) reducing the compound of formula (2) to give a compound of formula (3):




- c) coupling the compound of formula (3) with a compound of formula (4):



- 25 to give a compound of formula (5):




d) when W represents -OP<sup>2</sup>, deprotecting and then oxidising the compound of formula (5) to give a compound of formula (6):




5 and

e) subjecting the compound of formula (5) when W represents =O, or compound of formula (6) to ring-opening, and removal of any remaining protecting groups, to give a compound of formula (7):




10

2. A process for the preparation of a compound of formula (2):




which comprises cyanating a compound of formula (1):



15


wherein Y represents a halo group, preferably Cl or Br; P<sup>1</sup> represents a protecting group, and W represents =O or -OP<sup>2</sup>, in which P<sup>2</sup> represents a protecting group.

3. A process for the preparation of a compound of formula (3):




20

which comprises reduction of a compound of formula (2):




wherein  $P^1$  represents a protecting group, and  $W$  represents  $=O$  or  $-OP^2$ , in which  $P^2$  represents a protecting group.


- 5 4. A process for the preparation of a compound of formula (5):



which comprises coupling the compound of formula (3):



- 10 with a compound of formula (4):



wherein

$R^1$  represents an alkyl group, such as a  $C_{1-6}$  alkyl group, and preferably an isopropyl group;

- 15  $R^2$  represents an aryl group, preferably a phenyl group;


$R^3$  represents an aryl group, preferably a 4-fluorophenyl group;

$X$  a group of formula  $-COZ$ , wherein  $Z$  represents  $-OR^4$ , in which  $R^4$  represents an alkyl, preferably a methyl or ethyl, group, or  $-NR^5R^6$ , wherein  $R^5$  and  $R^6$  each independently represent H, alkyl, or aryl, and preferably  $R^5$  is H and  $R^6$  is phenyl;

- 20  $P^1$  represents a protecting group; and


$W$  represents  $=O$  or  $-OP^2$ , in which  $P^2$  represents a protecting group.

5. A compound of formula (2):




wherein  $P^1$  represents a protecting group, and W represents  $=O$  or  $-OP^2$ , in which  $P^2$  represents a protecting group.

6. A compound of formula (3):



wherein  $P^1$  represents a protecting group, and W represents  $=O$  or  $-OP^2$ , in which  $P^2$  represents a protecting group.

7. A compound of formula (5):



wherein

$R^1$  represents an alkyl group, such as a  $C_{1-6}$  alkyl group, and preferably an isopropyl group;

$R^2$  represents an aryl group, preferably a phenyl group;

$R^3$  represents an aryl group, preferably a 4-fluorophenyl group;

X a group of formula  $-COZ$ , wherein Z represents  $-OR^4$ , in which  $R^4$  represents an alkyl, preferably a methyl or ethyl, group, or  $-NR^5R^6$ , wherein  $R^5$  and  $R^6$  each independently represent H, alkyl, or aryl, and preferably  $R^5$  is H and  $R^6$  is phenyl;

$P^1$  represents a protecting group; and

W represents  $-OP^2$ , in which  $P^2$  represents a protecting group.

PCT/GB2004/003206

